

Scaling Properties of the Two-Chain Model

A.A.Nersesyan, A.Luther and F.V.Kusmartsev

February 1, 2008

*Institute of Theoretical Physics, Chalmers University of Technology,
S – 41296, Goteborg, Sweden*¹

and

*Institute of Physics, Georgian Academy of Sciences,
Tamarashvili 6, 380077, Tbilisi, Georgia*

NORDITA, Blegdamsvej 17, DK – 2100 Copenhagen 0, Denmark

L.D.Landau Institute for Theoretical Physics

Moscow, 117940, GSP – 1, Kosygin 2, V – 334, Russia

and

Department of Theoretical Physics,

Oulu University, Linnanmaa, 90570, Oulu, Finland

Abstract

Scaling properties of a self-dual field-theoretical model, describing two weakly coupled spinless Luttinger chains, are studied. A crossover to a sine-Gordon massive phase, with strongly developed two-particle interchain correlations, is described. It is argued that, in a wide range of the in-chain interaction, renormalization of the interchain hopping amplitude is determined by the Luttinger liquid effects.

¹Mailing address

e-mail: ners@fy.chalmers.se

1. Anderson suggested that in an array of weakly coupled chains a single-fermion interchain hopping ($t_{\perp}$) may be irrelevant (Anderson's confinement) [1], implying the possibility of a Luttinger liquid (LL) behavior of higher-dimensional ($d > 1$) strongly correlated electron systems [2]. The issue of Anderson's confinement, which is indeed a rather complicated problem even for chains with spinless fermions, has attracted much interest [1, 3 - 8].

Two opposite tendencies, associated with single-particle and two-particle correlations, characterize the low-energy behavior of a multi-chain system. On increasing the in-chain interaction g , the renormalized single particle hopping amplitude, being a measure of coherent delocalization of the particles between the chains, gets gradually suppressed due to infrared catastrophe [6]. In the spinful case this suppression is also contributed by the charge-spin separation [1]. At least in continuum models with a linear spectrum, such a tendency eventually leads to deconfinement-confinement transitions when the interaction becomes large enough [3,5,6]. On the other hand, depending on the sign of g , either particle-hole (ph) or particle-particle (pp) interchain hoppings are generated in the second order in $t_{\perp}$ [9,10]. These two-particle processes, with amplitudes increasing with $|g|$, inevitably drive the system away from the LL fixed point to strong-coupling, massive phases characterized by density-wave or pairing fluctuations, strongly correlated in the transverse direction [4 - 6].

Thus, the problem one faces when studying scaling properties of the two-chain system is the interplay between single-particle interchain hopping and two-particle correlations, described by two relevant operators with different critical dimensions. Understanding of this interplay is especially important at such values of $|g|$, when the two-particle hopping becomes the most relevant perturbation. It is not *a priori* clear what is the role of a large mass gap, generated by two-particle processes, in the renormalization of $t_{\perp}$. It is well known that such a mass gap could give rise to a finite threshold for a perturbation having the same scaling dimension as that of $t_{\perp}$. Such a situation occurs in 1d Fermi systems with attractive interaction in a magnetic field [11], or in "commensurate-incommensurate" transitions [12], described by the sine-Gordon model with a finite density of topological charge. Therefore, a question arises: Is it the LL "*infrared catastrophe*" or the two-particle mass gap which mostly determines renormalization of $t_{\perp}$? Is there any finite threshold for $t_{\perp}$, or not?

Motivated by these questions, we analyze scaling properties of a self-dual field-theoretical model, recently proposed [6] to describe two weakly coupled spinless

Luttinger chains. The model naturally incorporates both the development of single-fermion confinement and interchain pair coherence. This was demonstrated in [6] by using an equivalent representation of the theory in terms of a 2d Coulomb gas of charge-monopole composites. We show that, on decreasing the energy scale, the two-chain system scales down to a non-Luttinger, massive phase with strong interchain correlations, effectively described by a sine-Gordon model. At weak interaction, when $t_\perp$ is the most relevant perturbation, such a crossover occurs via formation of intermediate two-band LL, with a finite Zeeman-like interband splitting. Among possible ground states of the two-chain system, we indicate the orbital aniferromagnet phase, previously discussed by two of us [13]. Specifically, this type of ordering occurs due to interchain forward scattering.

At larger interaction, when the two-particle hopping dominates over the single-fermion one, the two-band LL regime is absent. Yet, we find no threshold for $t_\perp$ and argue that $t_\perp$ is mostly renormalized by pure LL screening effects. This conclusion is supported by using a relationship with spin-chain models.

2. We consider the following model of two spinless Luttinger chains with a weak single-particle interchain hopping:

$$H = \sum_{\mu} \int dx [-iv_F(\psi_{1\mu}^+ \partial_x \psi_{1\mu} - \psi_{2\mu}^+ \partial_x \psi_{2\mu}) + \pi v_F(g\rho_{1\mu}\rho_{2\mu} + g'\rho_{1\mu}\rho_{2,-\mu})] \\ + t_\perp \sum_{\mu} \int dx (\psi_{1\mu}^+ \psi_{1,-\mu} + \psi_{2\mu}^+ \psi_{2,-\mu}) \quad (1)$$

where v_F is the Fermi velocity, $\mu = \pm 1$ is the chain index, $\psi_{1\mu}$ and $\psi_{2\mu}$ are the Fermi fields for right-movers and left-movers, respectively, and $\rho_{j\mu}(x) = : \psi_{j\mu}^+(x)\psi_{j\mu}(x) :$ are the density operators. g and g' are dimensionless coupling constants characterizing the in-chain and interchain forward scattering, respectively. Using the standard Abelian bosonization, we transform the original problem of Fermi fields to a purely bosonic one, $H_B = H_\rho^0 + H_\sigma$ where H_ρ^0 is a Tomonaga-Luttinger Hamiltonian describing gapless excitations of the total (or "charge") density, $\rho_j = (1/\sqrt{2}) \sum_{\mu} \rho_{j\mu}$, which decouple from the rest of the spectrum. All nontrivial effects caused by the interchain hopping $t_\perp$ are incorporated in the "spin" part of the Hamiltonian $H_\sigma = H_\sigma^0 + H_\perp$ which deals with relative degrees of freedom, $\sigma_j = (1/\sqrt{2}) \sum_{\mu} \mu \rho_{j\mu}$. The coupling constants emerging in the ρ - and σ -channels are, respectively, $g_{\rho,\sigma} = g \pm g'$. As shown in [6], H_σ is given by the following field-theoretical model

$$H_\sigma = \int dx [\frac{u}{2} (P^2 + (\partial_x \phi)^2) + \frac{2t_\perp}{\pi\alpha} \cos(\frac{1}{2}\gamma\phi) \cos(\frac{1}{2}\tilde{\gamma}\tilde{\phi})] \quad (2)$$

Here $\phi(x)$ and $P(x)$ are a scalar field and its conjugate momentum, respectively, with a canonical commutation relation $[\phi(x, t), P(x', t)] = i\delta(x - x')$, $u = v_F(1 - \frac{1}{4}g_\sigma^2)^{1/2}$ is the renormalized velocity, $\tilde{\phi}(x)$ is a field dual to $\phi(x)$, defined as $\partial_x \tilde{\phi}(x) = P(x)$, and

$$\frac{\gamma^2}{8\pi} = \frac{8\pi}{\tilde{\gamma}} \equiv K = (1 - g_\sigma/2)^{1/2}(1 + g_\sigma/2)^{-1/2} \quad (3)$$

The fields ϕ and $\tilde{\phi}$ are related to the "spin" density and current: $\sigma_1 + \sigma_2 = (\gamma/\sqrt{8\pi})\partial_x \phi$, $\sigma_1 - \sigma_2 = -(\tilde{\gamma}/\sqrt{8\pi})\partial_x \tilde{\phi}$. The model (2) is self-dual: H_σ is invariant under transformations $\phi \leftrightarrow \tilde{\phi}$ ($g_\sigma \rightarrow -g_\sigma$), $\gamma \leftrightarrow \tilde{\gamma}$.

The perturbation term in (2) has critical dimension $\Delta = \frac{1}{2}(K + \tilde{K})$ [3] and, hence, is relevant when $\Delta < 2$. This inequality corresponds to the interval $K_- < K < K_+$, where the points $K_\pm = 2 \pm \sqrt{3}$ mark the boundaries between the confinement and deconfinement phases [6]. As follows from the Coulomb gas representation of the model (2) [6], the single-particle confinement originates from pairing of the charge-monopole composites with zero total "electric" and "magnetic" charges. On the other hand, binding of composites in pairs with total "magnetic" or "electric" charge ± 2 is described by two operators

$$O_{ph} = (\psi_{1\mu}^+ \psi_{1,-\mu})(\psi_{2,-\mu}^+ \psi_{2\mu}) \rightarrow -\frac{1}{2(\pi\alpha)^2} \cos \gamma \phi, \quad (4)$$

$$O_{pp} = (\psi_{1\mu}^+ \psi_{1,-\mu})(\psi_{2\mu}^+ \psi_{2,-\mu}) \rightarrow -\frac{1}{2(\pi\alpha)^2} \cos \tilde{\gamma} \tilde{\phi} \quad (5)$$

associated with interchain particle-hole (ph) and particle-particle (pp) hoppings, respectively. These two-particle processes, although absent in the initial Hamiltonian (2), are generated upon renormalization. The critical dimensions of operators (4) and (5) are, respectively, $2K$ and $1/2K$. So, at any $K \neq 1$ ($g_\sigma \neq 0$), one of these operators is always relevant. Therefore, the two-particle hopping processes must be included into renormalization scheme [10,14].

At the l -th step of the renormalization procedure (l can be identified as a continuous logarithmic variable $l = \ln(\alpha/\alpha_0)$ or $l = \ln(\Lambda/|\omega|)$), the Hamiltonian density takes the form

$$H_l(x) = \frac{u_l}{2} \left(P^2 + (\partial_x \phi)^2 \right) + \frac{4u_l \tau_l}{\alpha^2} \cos\left(\frac{1}{2}\gamma_l \phi\right) \cos\left(\frac{1}{2}\tilde{\gamma}_l \tilde{\phi}\right) - \frac{2\pi u_l}{(2\pi\alpha)^2} \left(G_l \cos(\gamma_l \phi) + \tilde{G}_l \cos(\tilde{\gamma}_l \tilde{\phi}) \right) \quad (6)$$

where $\gamma_l = 8\pi/\tilde{\gamma}_l$ is given by formula (3), in which g_σ is replaced by z_l ; $\tau_l = t_\perp(l)\alpha/2\pi u_l$ is a renormalized dimensionless single-fermion hopping amplitude.

The parameters of the Hamiltonian H_l are determined from the following renormalization-group equations ²

$$\tau_l' = (2 - \Delta_l)\tau_l \quad (7)$$

$$G_l' = 2(1 - K_l)G_l + (K_l - \tilde{K}_l)\tau_l^2 \quad (8)$$

$$\tilde{G}_l' = 2(1 - \tilde{K}_l)\tilde{G}_l + (\tilde{K}_l - K_l)\tau_l^2 \quad (9)$$

$$(\ln K)' = \frac{1}{2}(\tilde{K}_l\tilde{G}_l^2 - K_lG_l^2) \quad (10)$$

with initial conditions $u_0 = v_F$, $K_0 = K$, $G_0 = \tilde{G}_0 = 0$, and $\tilde{K}_l K_l = 1$. The upper prime indicates the derivative $\partial/\partial l$. Equations (8) and (9) were given by Yakovenko [14]. However, he did not consider renormalization of the coupling constant g_σ , Eq.(10), which, as will be shown below, is crucial for a weak interaction, when a crossover from a two-chain Luttinger liquid to a strong-coupling low-energy regime is developed.

3. We consider first the case of a weak interaction, when one can set

$$K_l \simeq 1 - \frac{1}{2}z_l, \quad \tilde{K}_l \simeq 1 + \frac{1}{2}z_l, \quad \Delta_l = 1 + 0(z_l^2). \quad (11)$$

Solving Eq.(7), one determines the value of the variable $l_0 = \ln(1/\tau_0) = \ln(\Lambda/\omega_0)$, at which τ_l becomes ~ 1 . This introduces a new energy scale to the problem - a Zeeman-like splitting of two degenerate bands: $\omega_0 \simeq \Lambda\tau_0 \sim t_\perp$.

In the region $l \leq l_0$, or $\omega_0 < |\omega| < \Lambda$, z_l is not renormalized (up to second-order corrections in g_σ), $z(l_0) = g_\sigma$, while the charges G_l and $\tilde{G}_l$ remain small:

$$G_l = -\tilde{G}_l = -\frac{1}{2}g_\sigma\tau_0^2(e^{2l} - e^{g_l}) \simeq -\frac{1}{2}g_\sigma\tau_l^2 \quad (12)$$

To solve the renormalization-group equations in the low-energy region $l \geq l_0$, we make use of a typical two-cutoff scaling prescription: understanding that at $l \sim l_0$ renormalization of $t_\perp(l)$ is stopped, we choose the normalization condition $\tau_{l_0} = 1$ and drop the τ^2 -terms in r.h.s. of Eqs.(8) and (9). Then we arrive at the following set of equations in the region $l \geq l_0$:

$$z' = G^2 - \tilde{G}^2, \quad G' = zG, \quad \tilde{G}' = -z\tilde{G} \quad (13)$$

²There is one more equation, $\zeta_l' = -4\pi^2(\Delta_l + 1)\tau_l^2$, which describes renormalization of the velocity: $u_l \rightarrow u_l(1 - \zeta_l)$. This renormalization reduces to rescaling of K and $\tilde{K}$: $K \rightarrow K(1 - \zeta^2)^{-1/2}$, $\tilde{K} \rightarrow \tilde{K}(1 - \zeta^2)^{1/2}$, which will be assumed everywhere below.

with the boundary conditions $z(l_0) = g_\sigma$, $G(l_0) = -\tilde{G}(l_0) = -\frac{1}{2}g_\sigma$.

Eqs.(13) describe a Z_4 -symmetric model. Its 2d classical version is a planar XY model in the presence of 4-fold degenerate symmetry breaking field [15,16]. Such a model is also equivalent to a version of the backscattering model of 1d fermions [17] extended to include spin-nonconserving processes [18]. We recognize a field-theoretical version of this model in (6) when $t_\perp$ is dropped. At $l = l_0$ this model reads

$$H_{l_0} = \frac{v_F}{2} \left(P^2 + (\partial_x \phi)^2 \right) + \frac{\pi v_F g_\sigma}{(2\pi\alpha)^2} (\cos \gamma \phi - \cos \tilde{\gamma} \tilde{\phi}) \quad (14)$$

Note that, in Eq.(14) amplitudes of the cosines coincide. This reflects the "hidden" $U(1)$ symmetry of the model (14) which brings it to the sine-Gordon (SG) universality class, corresponding to a critical plane of the Z_4 -model [19]. This is directly seen from the solution of Eqs.(13). Introducing linear combinations $G_\pm = G \pm \tilde{G}$ and taking into account boundary conditions at $l = l_0$, one finds that, at all $l > l_0$, $z = -G_-$, thus reducing Eqs.(13) to a pair of Kosterlitz-Thouless equations for the SG model: $G'_+ = -G_-^2$, $G'_- = -G_+ G_-$. The conditions $G_+(l_0) = 0$, $G_-(l_0) = -g_\sigma$ define a scaling trajectory with the initial point lying on the vertical axis of the standard Kosterlitz-Thouless phase plane, $(G_+, |G_-|)$. Therefore, for any sign of g_σ , a strong-coupling regime develops in the infrared limit. The dynamically generated mass gap

$$M \simeq t_\perp \exp(-\pi/2|g_\sigma|) \ll t_\perp \quad (15)$$

signals the onset of strong interchain two-particle correlations. The fact that the mass does not depends on the sign of g_σ is related to self-dual symmetry of the model.

4. Let us consider our model in the new basis which is built up from symmetric and antisymmetric states $\Psi_{j,\pm} = (\pm\psi_{j,+} + \psi_{j,-})/\sqrt{2}$. In the absence of the interchain hopping, two bands associated with these states are degenerate. The degeneracy is removed by interchain hopping, and the resulting Zeeman-like splitting of the two bands is a measure of coherent delocalization of fermions between the chains.

When the chain index is formally treated as a spin-1/2 variable, the interchain hopping term of the Hamiltonian turns to Zeeman interaction with a magnetic field $2t_\perp$ along the x -axis. In the new basis of states, which is obtained by a $\pi/2$ -

rotation about the y -axis in "spin" space, the magnetic field is oriented along the z -axis, and the splitting of the bands is determined by "magnetization" $< S^z >$.

The $\pi/2$ - rotation of the basis has no effect on the "charge" degrees of freedom. However, it produces the following changes in the σ -channel:

$$O_{ph,pp} \rightarrow \sigma_1 \sigma_2 \pm \frac{1}{2}(O_{ph} - O_{pp}), \quad \sigma_1 \sigma_2 \rightarrow \frac{1}{2}(O_{ph} + O_{pp}) \quad (16)$$

Quantities in the r.h.s. of (16) now have a different meaning: σ_j is a difference between the particle densities in each of the bands, while the operators O_{ph} and O_{pp} describe *interband* pair-hopping processes

Making use of (16), we transform the two-chain Hamiltonian (6) to a *two-band* one:

$$H'_l = \frac{u_l}{2} \left(\Pi^2 + (\partial_x \chi)^2 \right) + \frac{\beta_l t_l}{2\pi} \partial_x \chi + \frac{2\pi u_l}{(2\pi\alpha)^2} \left(z_\perp(l) \cos(\beta_l \chi) + z_f(l) \cos(\tilde{\beta}_l \tilde{\chi}) \right) \quad (17)$$

where

$$\begin{aligned} \frac{\beta_l^2}{8\pi} &= \frac{8\pi}{\tilde{\beta}_l^2} = \left(\frac{1 - \frac{z_\parallel(l)}{2}}{1 + \frac{z_\parallel(l)}{2}} \right)^{1/2}, \quad z_\parallel(l) = G_l + \tilde{G}_l \\ z_\perp(l) &= \frac{1}{2}(z_l + G_l - \tilde{G}_l), \quad z_f(l) = \frac{1}{2}(z_l - G_l + \tilde{G}_l) \end{aligned} \quad (18)$$

with initial conditions

$$z_\parallel(0) = 0, \quad z_\perp(0) = z_f(0) = \frac{1}{2}g. \quad (19)$$

In Eq. (17) we recognize the above mentioned U(1) symmetric 1d Fermi-gas model with backscattering ($z_\perp$) and spin-nonconserving processes (z_f) in a magnetic field along the z -axis [19]. At $l \leq l_0$ the effective couplings equal

$$z_\parallel(l) = 0, \quad z_\perp(l) = \frac{1}{2}g_\sigma(1 - \tau_l^2), \quad z_f(l) = \frac{1}{2}g_\sigma(1 + \tau_l^2). \quad (20)$$

One sees that, at energies $|\omega| \sim t_\perp$ ($l \sim l_0$), splitting of the two bands (τ_l) tends to suppress backscattering (or interband ph transitions, $z_\perp(l)$), which is due to nonconservation of the total momentum. The value of l_0 in the two-cutoff scaling is determined by condition $z_\perp(l) = 0$.

The suppression of the $z_\perp$ -term breaks the tiny balance between the amplitudes of the two cosines, which preserves a Tomonaga-Luttinger weak-coupling behavior at $t_\perp = 0$, and results in a crossover to a SG model

$$H'_{l_0} = \frac{v_F}{2} \left(\tilde{\Pi}^2 + (\partial_x \tilde{\chi})^2 \right) + \sqrt{\frac{2}{\pi}} t_{l_0} \tilde{\Pi} - \frac{2\pi v_F}{(2\pi\alpha)^2} g_\sigma \cos(\tilde{\beta}_0 \tilde{\chi}), \quad (21)$$

where $\tilde{\beta}_0 = \beta_0 = \sqrt{8\pi}$. This model is written in terms of the dual field $\tilde{\chi}$, using the relationship $\partial_x \chi = \tilde{\Pi}$. Notice that t_{l_0} is coupled to the momentum of the sine-Gordon field $\tilde{\chi}$. Then, by Galilean invariance, the term linear in $\tilde{\Pi}$ is eliminated by a shift $\tilde{\Pi} = \tilde{\Pi}' - (\sqrt{2}t_{l_0}/\sqrt{\pi}v_F)$ and does not influence the scaling properties of the system when the energy is further decreased in the region $l > l_0$. The remaining SG model is precisely the rotated version of H_{l_0} in (14). Since $\langle \tilde{\Pi}' \rangle = 0$, a response to the $t_{\perp}$ -perturbation, being just the magnetization of the "rotated" (the two-band) model, equals

$$\langle S^z \rangle_{l=l_0} = -\sqrt{\frac{2}{\pi}} \langle \partial_x \chi \rangle_{l=l_0} = -\sqrt{\frac{2}{\pi}} \langle \tilde{\Pi} \rangle_{l=l_0} = \frac{2t_{l_0}}{\pi v_F} \quad (22)$$

One has to take into account scale transformation of the coordinate x : $(\partial_x \chi)_{l=0} = e^{-l_0} (\partial_x \chi)_{l=l_0}$. Then one finds that t_{l_0} should be changed by $t_{\perp}$ in (22) when true response $\langle S^z \rangle_{l=0}$ is considered.

Since the $0(g^2)$ -correction to Δ was omitted, the obtained two-band splitting coincides with that for the noninteracting fermions. It can be easily checked that keeping such a correction is still within the main accuracy to which the above renormalization treatment of the case $|g_{\sigma}| \ll 1$ is valid. Then we obtain

$$\langle S^z \rangle = \frac{2t_{\perp}}{\pi v_F} \left(1 - \frac{1}{8} g_{\sigma}^2 \ln \frac{1}{\tau_0} + \dots \right). \quad (23)$$

The strong-coupling regime developing in the low-energy region ($|\omega| \leq \omega_0$) is related to ordering of the *dual* field $\tilde{\chi}$. In terms of the original model (6), this includes both possibilities for ordering of the field ϕ at $g_{\sigma} > 0$ (*ph* hopping) and the dual field $\tilde{\phi}$ at $g_{\sigma} < 0$ (*pp* hopping). The field χ remains disordered and, therefore, is free to adjust itself to a finite "magnetic field" $2t_{\perp}$, thus acquiring a nonzero expectation value $\langle \partial_x \chi \rangle$.

5. Let us clarify the symmetry of the ground state of the system, characterized by ordering of the field $\tilde{\chi}$. As follows from (21), its vacuum expectation value depends on the sign of g_{σ} : $\langle \tilde{\chi} \rangle = 0$ at $g_{\sigma} > 0$, and $\langle \tilde{\chi} \rangle = \pi/2\tilde{\beta}_0$ at $g_{\sigma} < 0$. Therefore, when bosonizing various bilinears in Fermi fields to construct order parameters in a $\rho - \sigma$ factorized form, only those operators proportional to $\cos(\tilde{\beta}_0 \tilde{\chi}/2)$ or $\sin(\tilde{\beta}_0 \tilde{\chi}/2)$ should be considered.

There are two pairs of competing states, with order parameters:

$$O_{CDW} = \sum_{\sigma} \Psi_{1\sigma}^+ \Psi_{2,-\sigma} = \sum_{\mu} \mu \psi_{1\mu}^+ \psi_{2\mu} \sim \exp(-i\sqrt{2\pi K_{\rho}} \chi_{\rho}) \cos \frac{1}{2} \tilde{\beta}_0 \tilde{\chi}, \quad (24)$$

$$O_{S1} = \sum_{\sigma} \Psi_{1\sigma} \Psi_{2\sigma} = \sum_{\mu} \psi_{1\mu} \psi_{2\mu} \sim \exp(-i\sqrt{2\pi\tilde{K}_{\rho}\tilde{\chi}_{\rho}}) \cos \frac{1}{2}\tilde{\beta}_0\tilde{\chi}, \quad (25)$$

and

$$O_{S2} = i \sum_{\sigma} \sigma \Psi_{1\sigma} \Psi_{2\sigma} = i \sum_{\mu} \psi_{1\mu} \psi_{2,-\mu} \sim \exp(-i\sqrt{2\pi\tilde{K}_{\rho}\tilde{\chi}_{\rho}}) \sin \frac{1}{2}\tilde{\beta}_0\tilde{\chi}, \quad (26)$$

$$O_{OAF} = i \sum_{\sigma} \sigma \Psi_{1\sigma}^+ \Psi_{2,-\sigma} = i \sum_{\mu} \mu \psi_{1\mu}^+ \psi_{2,-\mu} \sim \exp(-i\sqrt{2\pi\tilde{K}_{\rho}\tilde{\chi}_{\rho}}) \sin \frac{1}{2}\tilde{\beta}_0\tilde{\chi} \quad (27)$$

Here the field χ_{ρ} and its dual field $\tilde{\chi}_{\rho}$ describe gapless fluctuations in the ρ -channel, while $K_{\rho} \simeq 1 - \frac{1}{2}g_{\rho}$ and $\tilde{K}_{\rho} \simeq 1 + \frac{1}{2}g_{\rho}$ ($|g_{\rho}| \ll 1$) are critical exponents which determine power-law asymptotics of the corresponding correlation functions. The order parameters (24) - (27) are given both in the two-band and two-chain representations. These are two 1d charge-density waves with "antiferromagnetic" interchain ordering (CDW), the in-chain (S1) and interchain (S2) Cooper pairings, and orbital antiferromagnetic state (OAF), characterized by nonzero local currents circulating in the two-chain system with $2k_F$ -periodicity [13]. The operator (27) actually describes only transverse (interchain) currents. The alternating longitudinal (in-chain) currents are easily shown to be proportional to $(\partial_x \chi) O_{OAF} \sim (t_{\perp}/t_{\parallel}) O_{OAF}$, as it should be due to the current conservation law. All operators, other than (24)-(27), characterize states with short-ranged correlations, exponentially decaying at distances $|x| > 1/M$.

The dominant ordering among CDW and S1 at $g_{\sigma} > 0$ and among S2 and OAF at $g_{\sigma} < 0$ is determined by the sign of g_{ρ} . The resulting phase diagram is shown in Fig.1. It reflects the dual symmetry of the model: States OAF and S2 are dual to CDW and S1, respectively, and transform to each other under $g_{\sigma} \rightarrow -g_{\sigma}$. Notice the important role of interchain forward scattering g' which, in addition to already known states [10,5], opens new possibilities for ordering of the two-chain system, such as the OAF state [13]. More exotic phases, *e.g.* spin nematic, are expected in the case of spin-1/2 particles.

6. At weak interaction the mass gap M , Eq.(15), characterizing strong interchain two-particle correlations, is exponentially small compared to the interband splitting ω_0 . The crossover from a two-chain LL ($\omega_0 \ll |\omega| \ll \Lambda$) to a strong-coupling, quasi-ordered phase ($|\omega| \leq M$) occurs via formation of a two-band LL ($M \ll |\omega| \ll \omega_0$). On increasing g_{σ} , the difference between ω_0 and M decreases, and at large enough interaction the two-band LL regime no longer exists. In this case, one of the two-particle hopping amplitudes, G_l or $\tilde{G}_l$, goes to strong

coupling at such values of l , when τ_l is still small [14]. Questions which naturally arise, are: Can a small Zeeman-like splitting of the two bands survive the presence of a large mass gap M ? Is the mechanism of suppression of the single-particle interchain hopping, caused by LL effects (infrared catastrophe), essentially modified by strongly developed pair-hopping processes ? These questions cannot be answered by a simple comparison of rates of increase of G_l (or $\tilde{G}_l$) and τ_l^2 [14], since such an oversimplified treatment neglects different physical origin of the two relevant operators.

Let us assume for definiteness that $g_\sigma > 0$) and consider the case $K_0 < \sqrt{2}-1$, when the rate of increase of G_l exceeds that of τ_l^2 . Neglecting renormalization of K , unimportant at large interaction, from Eqs.(7) - (9) we obtain

$$G_l \simeq -\frac{K_0 - \tilde{K}_0}{2 + K_0 - \tilde{K}_0} \tau_0^2 e^{2(1-K_0)l}, \quad \tilde{G}_l \simeq \frac{\tilde{K}_0 - K_0}{2 + \tilde{K}_0 - K_0} \tau_l^2 \quad (28)$$

We see that G_l is increasing in such a way, as if its initial value were $\sim \tau_0^2$. The mass gap is then estimated as

$$M \simeq \Lambda [C(K_0)\tau_0^2]^{\frac{1}{2(1-K_0)}} \quad (29)$$

where $C(K_0)$ is a positive (nonuniversal) number.

Note that $G_l < 0$. Moreover, after a few renormalizations, $|G_l| \gg \tilde{G}_l$. This allows to estimate parameters of the "rotated" Hamiltonian (17):

$$z_{\parallel}(l) \simeq -|G_l|, \quad z_{\perp}(l) \simeq \frac{1}{2}(g_l - |G_l|), \quad z_f(l) \simeq \frac{1}{2}(g_l + |G_l|) \quad (30)$$

Then we find that β_l^2 defined by (18) is larger than 8π and increases upon renormalization. This indicates irrelevance of $\cos \beta_l \chi$ and relevance of $\cos \tilde{\beta}_l \tilde{\chi}$ in (17). This is also in agreement with scaling behavior of the effective amplitudes $z_{\perp}(l)$ and $z_f(l)$: the former increases upon renormalization, while the latter decreases.

These observations allow us to conclude that, even in the case when the interaction is not small, the model (17) scales towards a strong-coupling regime characterized by ordering of the dual field $\tilde{\chi}$. Then, following arguments used in sec.4, we expect the average magnetization $\langle S^z \rangle$ to be basically determined by scaling equation (7); therefore the interband splitting is given by

$$\omega_0 \simeq \Lambda \tau_0^{\frac{1}{2-\Delta_0}} \quad (31)$$

Of course, such estimations are valid up to some nonuniversal renormalizations, inevitable when interaction is not small. Nevertheless, it seems quite convincing that a large mass gap, indicating strong interchain pair coherence, coexists

with a finite interband splitting. The relevance of the single-fermion hopping is then mostly determined by the Luttinger critical dimension of the $t_{\perp}$ -perturbation in the original two-chain Hamiltonian (2). Therefore the statement, made in [6] concerning the Kosterlitz-Thouless-like nature of confinement-deconfinement transitions, is basically correct.

The fact that the "rotated" model (17) always shows ordering of the dual field $\tilde{\chi}$ is related to self-duality of the original model (2), still preserved when pair hopping is taken into account. As a counter example, violating this property, one could consider a two-chain model with interchain backscattering,

$$g_b \psi_{1\mu}^+ \psi_{2,-\mu}^+ \psi_{1,-\mu} \psi_{2\mu} \sim g_b \cos(\gamma\phi)$$

which changes the initial condition for G_l : $G_0 = g_b$. Using the "rotated" version of the model, Eq.(17), and repeating the same analysis as that in sec.3 and 4, it can be easily shown that, under certain conditions ($g_b < 0$, $0 < g_{\sigma} < |g_b|$), the interchain backscattering drives the system to a new strong-coupling fixed point. The latter is characterized by irrelevance of the dual field $\tilde{\chi}$ and ordering of the field χ , which describes two, "ferromagnetically" correlated, charge-density waves along the chains. This type of ordering does tend to suppress the effective single-particle interchain hopping, since $t_{\perp}$ now is coupled to the density of topological charge of the SG field χ . Like in the attractive 1d Fermi system in a magnetic field [11], a finite interband splitting will be only nonzero, if $t_{\perp}$ exceeds the corresponding mass gap, with square-root singularities near the threshold. In such a situation, the confinement-deconfinement transition, taking place on increasing the amplitude $t_{\perp}$, is of different nature.

7. To support the above conclusion concerning the existence of a finite band splitting in the presence of a large mass gap, we use a relationship between the two-chain model (2) and quantum spin-1/2 chains. Consider an aniferromagnetic XY spin-1/2 chain in a homogeneous magnetic field in the basal plane (e.g. along the x-axis)

$$H = J \sum_{n=1}^N (S_n^x S_{n+1}^x + S_n^y S_{n+1}^y) + h \sum_{n=1}^N S_n^x, \quad (J > 0) \quad (32)$$

and assume that $|h| \ll J$. The last condition makes it possible to consider the continuum limit of the model (32). Using the Jordan-Wigner representation of the spin operators and the standard bosonization procedure [20], we arrive at the

following field theory:

$$H = \frac{v_F}{2} \int dx \left(P^2(x) + (\partial_x \phi)^2 \right) + \frac{h}{\sqrt{2\pi\alpha a}} \int dx \cos(\sqrt{4\pi}\phi(x)) \cos(\sqrt{\pi}\tilde{\phi}) \quad (33)$$

We see that Hamiltonian (33) is a special case of the model (2) corresponding to the value $K = 2$ or, by duality, $K = 1/2$. At these particular values of K the spin-chain model (32), with $h \sim t_\perp$, represents a lattice version of the two-chain model.

Although the exact solution of the spin model (32) is unknown, we expect a finite homogeneous magnetization along the x-axis to exist in the ground state, which is equivalent to a finite delocalization of fermions between the chains. At $h = 0$ the ground state of the XY-model is disordered due to continuous $U(1)$ symmetry, and the excitation spectrum is gapless. The magnetic field, breaking $U(1)$ down to Z_2 , will lead to the appearance of a finite magnetization along the x-axis and to the opening of a gap, related to antiferromagnetic long-range ordering of the spins along the y-direction. The spin order parameter is given by the staggered magnetization $(-1)^n < S_n^y >$, which, in the continuum limit, reduces to $< \sin(\sqrt{\pi}\tilde{\phi}(x)) >$ at $K = 2$, or $< \sin(\sqrt{\pi}\phi(x)) >$ at $K = 1/2$, the latter two averages being the order parameters of the model (6), related to the development of the ph or pp pair coherence in the two-chain system.

A more detailed study of the correspondence between the two-chain problem and quantum spin chains will be presented elsewhere.

One of the authors (A.N.) acknowledges financial support from Chalmers University of Technology. He also would like to thank H.Johannesson, S.Kivelson, Yu Lu, T.Einarsson and S.Ostlund for helpful discussions on the two-chain problem. F.K. is grateful to C.Bourbonnais for many fruitful discussions. Two of us (F.K. and A.N.) acknowledge the generous hospitality of Nordita, where part of this work was done.

References

- [1] P.W.Anderson, Phys.Rev.Lett. 67 (1991) 3844.
- [2] P.W.Anderson, Phys.Rev.Lett. 64 (1990) 1839; 65 (1990) 2306.
- [3] X.G.Wen, Phys.Rev. B 42 (1990) 6623.
- [4] H.J.Schulz, Int.J.Mod.Phys. B 5 (1991) 57.
- [5] M.Fabrizio, A.Parola and E.Tosatti, Phys.Rev. B 46 (1992) 3159.
- [6] F.V.Kusmartsev, A.Luther and A.Nersesyan, JETP Lett. 55 (1992) 692.
- [7] C.Castellani, C.Di Castro and W.Metzner, Phys.Rev.Lett. 69 (1992) 1703.
- [8] A.Finkel'shtein and A.I.Larkin, to be published.
- [9] S.A.Brazovskii and V.M.Yakovenko, J.Physique Lett.(France) 46 (1985) L-111; Sov.Phys.JETP 62 (1985) 1340.
- [10] C.Bourbonnais and L.G.Caron, Int.J.Mod.Phys. B 5 (1991) 11033.
- [11] G.I.Japaridze and A.A.Nersesyan, J.Low Temp.Phys. 37 (1979) 95.
- [12] V.L.Pokrovsky and A.L.Talapov, Phys.Rev.Lett 42 (1979) 65.
- [13] A.A.Nersesyan and A.Luther, Nordita report, 1988; A.A.Nersesyan, Phys.Lett. A 153 (1991) 49.
- [14] V.M.Yakovenko, to be published.
- [15] J.V.Jose, L.P.Kadanoff, S.Kirkpatrick and D.R.Nelson, Phys.Rev. B 16 (1977) 1217.
- [16] P.B.Wiegmann, J.Phys. C11 (1978) 1583.
- [17] A.Luther and V.J.Emery, Phys.Rev.Lett. 33 (1974) 589.
- [18] T.T.Truong and K.D.Schotte, Phys.Rev.Lett. 47 (1981)
- [19] T.Giamarchi and H.J.Schulz, J.Phys. (Paris) 49 (1988) 819.
- [20] A.Luther and I.Peschel, Phys.Rev. B 12 (1975) 3908.

FIGURE CAPTION

Fig.1

The phase diagram on the plane (g, g') . g and g' are the in-chain and interchain forward scattering amplitudes, respectively.